

Preparation for Your Upcoming Colonoscopy

YOU WILL NEED:

Appointment Info

- SuTab® prescription filled by your pharmacy
- For coupon and info visit www.sutab.com

If you didn't collect your prep prescription shortly after we sent it to your pharmacy, kindly contact your pharmacy to have it prepared again.

Appointment into	Diabetes in structions (ii applicable)
DATE	Oral Medication Instructions
ARRIVAL TIME	Insulin Schedule
Endoscopy Center of Marin 1100 S. Eliseo Dr. Suite 3, Greenbrae Expect to be at the center 90 minutes	Blood Thinner Instructions (if applicable)
MarinHealth Medical Center 250 Bon Air Road, Greenbrae Expect to be at the hospital 2 hours	Continue blood thinner as prescribed HOLD for days
Novato Community Hospital 180 Rowland Way, Novato Expect to be at the hospital 3 hours	Weight Loss Prescriptions (if applicable) Weekly Dosing(Ozempic, Rybelsus, Wegovy, Trulicity, Mounjaro) Your last dose must be done 7 days before your procedure.
	Daily Dosing(Saxenda, Byetta) Do not take on the day of the procedure.
	Other instruction:

MarinHealth Gastroenterology | A UCSF Health Clinic

200 Tamal Plaza | Suite 200 | Corte Madera, CA 94925 7120 Redwood Blvd. Novato, CA 94945 415-925-6900 | MyMarinHealth.org/gastroclinic



Dighetes Instructions (if applicable)





- Confirm ride for the day of your procedure
- If you take blood thinners or weight loss medication, please review instructions for holding prescription.

Last chance to cancel/reschedule your procedure appointment. Please be aware that the Endoscopy Center of Marin may impose a late cancellation charge of up to \$200.



5 Days Before Your Colonoscopy

- Review thorough prep instructions and plan your meals
- Create a list of your medications and supplements to provide the facility. Include prescription name, dose, and directions
- Create a list of your allergies to provide the facility



3 Days Before Your Colonoscopy

- STOP eating foods with nuts, seeds, or whole grains. (fruits and vegetables with seeds, sesame, etc.)
- Avoid corn, beans, quinoa, and popcorn
- For more diet recommendations, please visit:
 MyMarinHealth.org/prep



2 Days Before Your Colonoscopy

- Increase your fluid intake, drink a minimum of 64 oz of clear liquids (approximately 8 glasses) (water preferred)
- Eat well-balanced meals, but continue avoiding nuts, seeds, and whole grains

TIPS

- To improve the taste try chilling laxative and/or using a straw to drink mixture
- If you experience nausea or vomiting, rest for 15 minutes, then try to resume prep
- Use unscented wipes instead of toilet paper.
- Use Vaseline or an ointment for diaper rash around the anus before and/or during prep to minimize irritation from passing bowel movements



"PREP DAY" — DOSE 1

- Have a low-residue breakfast BEFORE 10 am (white toast, eggs, yogurt, cream of wheat, smoothies without seeds)
- At 10 am begin a "Clear Liquid Diet." Clear liquids include water, juice without pulp, soda, black coffee/ tea (no milk), electrolyte drinks, Ensure CLEAR, Boost Breeze®, Jell-O, broth, popsicles
- At 6pm take the first dose of SuTab®

Take the tablets with water

Step 1: Open 1 bottle of 12 tablets

Step 2: Fill the provided container with 16 oz of water (up to the fill line). Swallow each tablet with a sip of water, and drink the entire amount of water over 15–20 minutes.

Drink additional water

Step 3: Approximately 1 hour after the last tablet is ingested, fill the provided container again with 16 oz of water (up to the fill line), and drink the entire amount over 30 minutes.

Step 4: Approximately 30 minutes after finishing the second container of water, fill the provided container with 16 oz of water (up to the fill line), and drink the entire amount over 30 minutes.

The Day of Your Colonoscopy

DOSE 2

- Continue to consume only clear liquids until after the colonoscopy
- The morning of the colonoscopy (at least 6 hours prior to your arrival time), open the second bottle of 12 tablets

Repeat Steps 2–4 from DOSE 1

- Please complete the bowel prep 3 hours before your colonoscopy arrival time. Please STOP drinking all liquids 3 hours before your colonoscopy arrival time.
- You should take your regular medications/ supplements at least three hours before your arrival time or after your colonoscopy.

BRING THE FOLLOWING:

Photo ID

- Medication List
- Insurance Card
- Allergy List